# NKMAXBio We support you, we believe in your research

# **Human PPME1 antibody**

Catalog Number: ATGA0449

## **PRODUCT INFORMATION**

## Catalog number

ATGA0449

#### Clone No.

AT29G3

## **Product type**

Monoclonal Antibody

#### UnitProt No.

Q9Y570

#### **NCBI Accession No.**

NP 057231

#### **Alternative Names**

Protein phosphatase methylesterase 1, FLJ22226, PME-1

## **PRODUCT SPECIFICATION**

## **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human PPME1 (1-386aa) purified from E. coli

## Isotype

IgG1 kappa

## **Purification Note**

By protein-A affinity chromatography

## **Application**

ELISA, WB

## Usage

The antibody has been tested by ELISA, Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

#### **Storage**



# NKMAXBIO We support you, we believe in your research

## **Human PPME1 antibody**

Catalog Number: ATGA0449

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

## **Description**


PPME1, also known protein phosphatase methylesterase-1 (PME1), catalyzes the demethylation and inactivation of protein phosphatase (PP2A), which is a multimeric phosphoserine/ threonine protein phosphatase associated with growth inhibition and cell cycle arrest. It can demethylate PP2A catalytic subunit in vitro and okadaic acid treatment is capable of inhibiting this reaction. It is conserved from yeast to human and contains a motif found in lipases having a catalytic triad activated serine as their active site nucleophile.

#### **General References**

Ogris E., et al. (1999) J. Biol. Chem. 274: 14382-14391. Gagnon S.N., et al. (2002) Biochem. J. 368: 263-271.

## **DATA**

## Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human PPME1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: K562 cell lysate Lane 3.: A549 cell lysate

