

Human Histone Deacetylase 2/HDAC2 antibody

Catalog Number: ATGA0510

PRODUCT INFORMATION

Catalog number

ATGA0510

Clone No.

AT3B1

Product type

Monoclonal Antibody

UnitProt No.

Q92769

NCBI Accession No.

NP_001518

Alternative Names

Histone deacetylase 2, HD2, RPD3, YAF1, KDAC2, Protein deacylase HDAC2

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human HDAC2 (1-488aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Human Histone Deacetylase 2/HDAC2 antibody

Catalog Number: ATGA0510

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


HDAC2 belongs to the histone deacetylase family that act via the formation of large multiprotein complexes and are responsible for the deacetylation of lysine residues on the N-terminal region of the core histones. It forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. It also plays an important role in transcriptional regulation, cell cycle progression and developmental events.

General References

Betz R. et al. (1998) Genomics. 52:245-6.
Taunton J. et al. (1996) Science 272:408-11.


DATA

Western blot analysis (WB)



The recombinant protein (50ng) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human HDAC2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: Recombinant Human HDAC2

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of HDAC2 in HeLa cells. The cell was stained with ATGA0510 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)

Human Histone Deacetylase 2/HDAC2 antibody

Catalog Number: ATGA0510

Flow cytometry analysis of HDAC2 in A549 cells. The cell was stained with ATGA0510 at 2-5ug for 1×10^6 cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (blue), cells without incubation with primary and secondary antibody was used as the negative control (black).

